

Aristocort

Aristocort - Aristocort (Triamcinolone) is used for treating a variety of medical conditions, including but not limited to, insufficiency of the adrenal cortex, inflammatory disorders, allergic disorders, and connective tissue disorders. It may also be used for other conditions as determined by your doctor.

Contents

- 1 Drug name
- 2 Drug Uses
- 3 How Taken
- 4 Drug Class and Mechanism
- 5 Missed Dose
- 6 Storage
- 7 Warnings Precautions
- 8 Possible Side Effects
- 9 More Information
- 10 Site:

Drug name

Aristocort (Triamcinolone)

Drug Uses

Aristocort (Triamcinolone) is used for treating a variety of medical conditions, including but not limited to, insufficiency of the adrenal cortex, inflammatory disorders, allergic disorders, and connective tissue disorders. It may also be used for other conditions as determined by your doctor.

How Taken

Use Aristocort as directed by your doctor. Check the label on the medicine for exact dosing instructions.

- Take Aristocort by mouth with food.
- To reduce side effects, take Aristocort around 8 a.m. in the morning.
- Do not take your medicine more often than directed.
- Do not stop taking Aristocort except on your prescriber's advice.

Drug Class and Mechanism

Aristocort is a corticosteroid. It works by reducing inflammation and modifying the body's immune system.

Missed Dose

If you miss a dose, take it as soon as you can. If it is almost time for your next dose, consult your prescriber or health care professional. You may need to miss a dose or take a double dose, depending on your condition and treatment. Do not take double or extra doses without advice.

Storage

Aristocort should be stored at room temperature, 59-86 degrees F (15-30 degrees C) and protected from heat, and direct light.

Warnings Precautions

Do not use Aristocort if:

- you are allergic to any ingredient in Aristocort;
- you have a fungal infection ;
- you are taking mifepristone .

Contact your doctor or health care provider right away if any of these apply to you. Before using Aristocort :

- Some medical conditions may interact with Aristocort . Tell your doctor or pharmacist if you have any medical conditions, especially if any of the following apply to you:
- if you are pregnant, planning to become pregnant, or are breast-feeding;
- if you are taking any prescription or nonprescription medicine, herbal preparation, or dietary supplement;
- if you have allergies to medicines, foods, or other substances;
- if you have or have been recently exposed to a bacterial, fungal, parasitic, ameba, or viral infection, especially measles, chickenpox, shingles, tuberculosis (TB), or herpes simplex of the eye;
- if you have been recently vaccinated, or have had a positive TB skin test;
- if you have heartburn, stomach pain, diarrhea, or intestinal problems (eg, obstructions, perforations, peptic ulcer, recent intestinal surgery);
- if you have a blood coagulation disorder, heart problems, diabetes mellitus, kidney problems, liver disease, an underactive thyroid, unstable joints, or osteoporosis, fractures, or have had a recent heart attack or joint surgery;
- you have unstable emotions, depression, or myasthenia gravis (muscle weakness).

Some medicines may interact with Aristocort . Tell your health care provider if you are taking any other medicines, especially any of the following:

- Barbiturates (eg, phenobarbital), carbamazepine, hydantoins (eg, phenytoin), or rifampin because they may decrease Aristocort 's effectiveness;
- Clarithromycin, oral contraceptives (birth control pills), or itraconazole because the risk of side effects, such as nervous system or adrenal gland problems, may be increased;
- Aspirin, mifepristone or ritodrine because the risk of their side effects may be increased by Aristocort.

Possible Side Effects

All medicines may cause side effects, but many people have no, or minor, side effects. Check with your doctor if any of these most common side effects persist or become bothersome:

- bone fractures; changes in appetite; changes in mood; difficulty sleeping; headache; indigestion; nervousness; stomach bloating; thin, fragile skin.

Seek medical attention right away if any of these severe side effects occur:

- severe allergic reactions (rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue); black, tarry stools; changes in menstrual periods; convulsions; fast or irregular heartbeat; prolonged sore throat, cold, or fever; psychotic behaviors; severe nausea or vomiting; swelling of feet or legs; unusual bruising or bleeding; unusual weight gain; vision problems; vomiting that looks like coffee grounds.

More Information

If you have any questions about Aristocort , please talk with your doctor, pharmacist, or other health care provider. Aristocort is to be used only by the patient for whom it is prescribed. Do not share it with other people. If your symptoms do not improve or if they become worse, check with your doctor.

Site:

[Aristocort](#)