

# Avapro

**Avapro** - Avapro is used for treating high blood pressure. It may be used alone or with other blood pressure medicines. It is also used for treating kidney disease in patients with type 2 diabetes.

## Contents

- 1 Drug name
- 2 Drug Uses
- 3 How Taken
- 4 Drug Class and Mechanism
- 5 Missed Dose
- 6 Storage
- 7 Warnings Precautions
- 8 Possible Side Effects
- 9 More Information
- 10 Site:

## Drug name

Avapro (Irbesartan)

## Drug Uses

Avapro is used for treating high blood pressure. It may be used alone or with other blood pressure medicines. It is also used for treating kidney disease in patients with type 2 diabetes.

## How Taken

Use Avapro as directed by your doctor.

- Take Avapro by mouth with or without food.
- Do not take 2 doses at once.

## Drug Class and Mechanism

Avapro is an angiotensin 2 receptor antagonist. It works by decreasing certain chemicals in the body, which cause blood vessels to narrow and sodium and fluids to be stored in the kidneys.

## Missed Dose

If you miss a dose of Avapro , take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

## Storage

Store Avapro between 39 and 86 degrees F (4 and 30 degrees C). Keep Avapro out of the reach of children and away from pets.

## Warnings Precautions

Do not use Avapro if:

- you are allergic to any ingredient in Avapro;
- you are pregnant and past your third month.

Contact your doctor or health care provider right away if any of these apply to you. Important:

- Avapro may cause drowsiness, dizziness, blurred vision, or lightheadedness. These effects may be worse if you take it with alcohol or certain medicines. Use Avapro with caution. Do not drive or perform other possibly unsafe tasks until you know how you react to it.
- Patients who take medicine for high blood pressure often feel tired or run down for a few weeks after starting treatment. Be sure to take your medicine even if you may not feel *normal*. Tell your doctor if you develop any new symptoms.
- Avapro may not work as well in black patients. They may also be at greater risk of side effects. Contact your doctor if your symptoms do not improve or if they become worse.
- Avapro should not be used in children younger than 6 years old; safety and effectiveness in these children have not been confirmed.
- Pregnancy and breast-feeding: Do not take Avapro after the third month of pregnancy. It may cause injury or death to the fetus. If you become pregnant, contact your doctor right away. It is not known if Avapro is found in breast milk. Do not breast-feed while taking Avapro.

## Possible Side Effects

Check with your doctor if any of these most common side effects persist or become bothersome:

- anxiety; chest pain; diarrhea; dizziness; flu; headache; heartburn/indigestion; muscle injury; tiredness; muscle pain; nausea; nervousness; runny nose; sinus abnormality; sore throat; stomach pain; swelling; upper respiratory tract infection; urinary tract infection; vomiting.

Seek medical attention right away if any of these severe side effects occur:

- severe allergic reactions (rash; hives; itching; difficulty breathing; tightness in the chest; swelling of the mouth, face, lips, or tongue); chest pain; fast heartbeat; fainting.

## More Information

Avapro is to be used only by the patient for whom it is prescribed. Do not share it with other people. If your symptoms do not improve or if they become worse, check with your doctor.

## Site:

Avapro